



The Tachykinins Substance P and Hemokinin-1 Favor the Generation of Human Memory Th17 Cells by Inducing IL-1 β , IL-23, and TNF-Like 1A Expression by Monocytes

Submitted by Emmanuel Lemoine on Thu, 03/26/2015 - 14:22

Titre	The Tachykinins Substance P and Hemokinin-1 Favor the Generation of Human Memory Th17 Cells by Inducing IL-1 β , IL-23, and TNF-Like 1A Expression by Monocytes
Type de publication	Article de revue
Auteur	Cunin, Pierre [1], Caillon, Antoine [2], Corvaisier, Murielle [3], Garo, Erwan [4], Scotet, Mari [5], Blanchard, Simon [6], Delneste, Yves [7], Jeannin, Pascale [8]
Editeur	American Association of Immunologists
Type	Article scientifique dans une revue à comité de lecture
Année	2011
Langue	Anglais
Date	2011/01/04
Numéro	7
Pagination	4175 - 4182
Volume	186
Titre de la revue	The Journal of Immunology
ISSN	1550-6606
Résumé en anglais	<p>The nervous system influences immune responses through the release of neural factors such as neuropeptides. Among them, the tachykinin substance P (SP) signals via the neurokinin 1 receptor (NK-1R), which is expressed by various immune cells. We thereby analyzed in this paper whether tachykinins may participate in human CD4⁺ Th cell polarization. We report that SP and hemokinin-1 (HK-1) upregulate IL-17A and IFN-γ production by human memory CD4⁺ T cells without affecting IL-4 and IL-10 production. SP and HK-1 switch non-Th17-committed CD4⁺ memory T cells into bona fide Th17 cells and Th1/Th17 cells. In contrast, SP and HK-1 do not modulate the polarization of naive CD4⁺ T cells. SP- and HK-1-induced Th17 cell generation is mediated through NK-1R and requires the presence of monocytes. SP and HK-1 trigger IL-1β, IL-6, and TNF-α production, upregulate IL-23 production, and enhance TNF-like 1A expression on monocyte surface. Neutralization experiments demonstrated that IL-1β, IL-23, and TNF-like 1A are involved in the SP- and HK-1-induced Th17 cell. The other members of the tachykinin family, neurokinins A and B, have no effect on the differentiation of naive and memory T cells. These results thereby show that SP and HK-1 are novel Th17 cell-inducing factors that may act locally on memory T cells to amplify inflammatory responses.</p>
URL de la notice	http://okina.univ-angers.fr/publications/ua9173 [9]
DOI	10.4049/jimmunol.1002535 [10]
Lien vers le document	http://dx.doi.org/10.4049/jimmunol.1002535 [10]

Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=16276](http://okina.univ-angers.fr/publications?f[author]=16276)
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=16305](http://okina.univ-angers.fr/publications?f[author]=16305)
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=7896](http://okina.univ-angers.fr/publications?f[author]=7896)
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=7900](http://okina.univ-angers.fr/publications?f[author]=7900)
- [5] [http://okina.univ-angers.fr/publications?f\[author\]=16183](http://okina.univ-angers.fr/publications?f[author]=16183)
- [6] <http://okina.univ-angers.fr/si.bla/publications>
- [7] [http://okina.univ-angers.fr/publications?f\[author\]=3873](http://okina.univ-angers.fr/publications?f[author]=3873)
- [8] <http://okina.univ-angers.fr/pascale.jeannin/publications>
- [9] <http://okina.univ-angers.fr/publications/ua9173>
- [10] <http://dx.doi.org/10.4049/jimmunol.1002535>

Publié sur *Okina* (<http://okina.univ-angers.fr>)